

# Periodic Table of the Finite Elements

	k=0	k=1	k=2	k=3		k=0	k=1	k=2	k=3		k=0	k=1	k=2	k=3		k=0	k=1	k=2	k=3	
n=1	r=1			$\mathcal{P}_r^{-1} \mathcal{A}^k$	<p>The shape function space for <math>\mathcal{P}_r \mathcal{A}^k</math> is <math>\mathcal{P}_{r-1} \mathcal{A}^k + \kappa \mathcal{P}_{r-1} \mathcal{A}^{k+1}</math>, where <math>\kappa</math> is the Koszul differential.<sup>7</sup> It includes the full polynomial space <math>\mathcal{P}_{r-1} \mathcal{A}^k</math>, is included in <math>\mathcal{P}_r \mathcal{A}^k</math>, and has dimension <math>\dim \mathcal{P}_r \mathcal{A}^k(\Delta_n) = \binom{r+n}{r+k} \binom{r+k-1}{k}</math>. The degrees of freedom are given on faces of dimension <math>d \geq k</math> by moments of the trace weighted by a full polynomial space: <math>u \mapsto \int_{\tau} (tr u) \wedge q, q \in \mathcal{P}_{r+k-d-1} \mathcal{A}^{d-k}(f)</math>. The spaces with constant degree <math>r</math> form a complex: <math>\mathcal{P}_r \mathcal{A}^0 \xrightarrow{d} \mathcal{P}_r \mathcal{A}^1 \xrightarrow{d} \dots \xrightarrow{d} \mathcal{P}_r \mathcal{A}^n</math>.</p>			$\mathcal{P}_r \mathcal{A}^k$	<p>The shape function space for <math>\mathcal{P}_r \mathcal{A}^k</math> consists of all differential <math>k</math>-forms with polynomial coefficients of degree at most <math>r</math>, and has dimension <math>\dim \mathcal{P}_r \mathcal{A}^k(\Delta_n) = \binom{r+n}{r+k} \binom{r+k}{k}</math>. The degrees of freedom are given on faces of dimension <math>d \geq k</math> by moments of the trace weighted by a <math>\mathcal{P}_r</math> space: <math>u \mapsto \int_{\tau} (tr u) \wedge q, q \in \mathcal{P}_{r+k-d} \mathcal{A}^{d-k}(f)</math>. The spaces with constant degree <math>r</math> form a complex: <math>\mathcal{P}_r \mathcal{A}^0 \xrightarrow{d} \mathcal{P}_r \mathcal{A}^1 \xrightarrow{d} \dots \xrightarrow{d} \mathcal{P}_r \mathcal{A}^n</math>.</p>			$\mathcal{Q}_r^{-1} \mathcal{A}^k$	<p>This family is constructed from the complex of 1-dimensional finite elements using a tensor product construction.<sup>10</sup> The shape function space on the unit cube <math>\square_n = I^n</math> is given by <math>\bigoplus_{\sigma \in \Sigma(k,n)} \bigotimes_{L=1}^n \mathcal{P}_{\sigma(L)}(I)</math> where <math>\Sigma(k,n)</math> denotes the increasing maps <math>\{1, \dots, k\} \rightarrow \{1, \dots, n\}</math>. Its dimension is <math>\dim \mathcal{Q}_r \mathcal{A}^k(\square_n) = \binom{n}{k} (r+1)^{n-k}</math>. The degrees of freedom are <math>u \mapsto \int_{\tau} (tr u) \wedge q, q \in \mathcal{Q}_{r-1} \mathcal{A}^{d-k}(f)</math>. The spaces with constant degree <math>r</math> form a complex: <math>\mathcal{Q}_r \mathcal{A}^0 \xrightarrow{d} \mathcal{Q}_r \mathcal{A}^1 \xrightarrow{d} \dots \xrightarrow{d} \mathcal{Q}_r \mathcal{A}^n</math>.</p>			$\mathcal{S}_r \mathcal{A}^k$	<p>The shape function space for <math>\mathcal{S}_r \mathcal{A}^k</math> is given by <math>\mathcal{P}_r \mathcal{A}^k \oplus \bigoplus_{\ell=0}^{k-1} \mathcal{H}_{r-\ell-1} \mathcal{A}^{k-1} \oplus \mathcal{H}_{r+k} \mathcal{A}^k</math>, where <math>\mathcal{H}_\ell \mathcal{A}^k</math> consists of homogeneous polynomial <math>k</math>-forms of degree <math>r</math> which are linear and undifferentiated in at least <math>\ell</math> variables.<sup>11</sup> Its dimension is <math>\dim \mathcal{S}_r \mathcal{A}^k(\square_n) = \sum_{\ell=0}^k 2^{d-\ell} \binom{r-d+2k}{\ell}</math>. The degrees of freedom are <math>u \mapsto \int_{\tau} (tr u) \wedge q, q \in \mathcal{P}_{r-2(d-k)} \mathcal{A}^{d-k}(f)</math>. The spaces with decreasing degree <math>r</math> form a complex: <math>\mathcal{S}_r \mathcal{A}^0 \xrightarrow{d} \mathcal{S}_{r-1} \mathcal{A}^1 \xrightarrow{d} \dots \xrightarrow{d} \mathcal{S}_{r-n} \mathcal{A}^n</math>.</p>			
	r=2																			
	r=3																			
n=2	r=1																			
	r=2																			
	r=3																			
n=3	r=1																			
	r=2																			
	r=3																			

**Legend**

Element with degrees of freedom (DOFs)

Symbol of element

Weight functions for DOFs

Dimension of element function space

Finite element exterior calculus notation

Element specification in FEM

	k=0	k=1	k=2	k=3
$\dim \mathcal{P}_r \mathcal{A}^k(\Delta_n) = \binom{r+n}{r+k} \binom{r+k-1}{k}$	2, 3, 4, 5, 6, 7, 8	3, 4, 5, 6, 7, 8	4, 5, 6, 7, 8	5, 6, 7, 8
$\dim \mathcal{P}_r \mathcal{A}^k(\Delta_n) = \binom{r+n}{r+k} \binom{r+k}{k}$	2, 3, 4, 5, 6, 7, 8	3, 4, 5, 6, 7, 8	4, 5, 6, 7, 8	5, 6, 7, 8
$\dim \mathcal{Q}_r \mathcal{A}^k(\square_n) = \binom{n}{k} (r+1)^{n-k}$	2, 3, 4, 5, 6, 7, 8	3, 4, 5, 6, 7, 8	4, 5, 6, 7, 8	5, 6, 7, 8
$\dim \mathcal{S}_r \mathcal{A}^k(\square_n) = \sum_{\ell=0}^k 2^{d-\ell} \binom{r-d+2k}{\ell}$	2, 3, 4, 5, 6, 7, 8	3, 4, 5, 6, 7, 8	4, 5, 6, 7, 8	5, 6, 7, 8

### Finite elements

The table presents the primary aspects of finite elements for the distribution of the fundamental operators of vector calculus: the gradient, curl, and divergence. A finite element space is a space of piecewise polynomial functions on a domain determined by (1) a mesh of the domain into polyhedral cells called elements, (2) a finite dimensional space of polynomial functions on each element called the shape functions, and (3) a consistent set of functionals on the shape functions of each element called degrees of freedom (DOFs), each DOF being associated to a generalized face of the element, and specifying a quantity which takes a single value at all elements sharing the face. The element diagrams depict the DOFs and their association to faces.

The spaces  $\mathcal{P}_r \mathcal{A}^k$  and  $\mathcal{P}_r \mathcal{A}^k$  depicted on the left half of the table are the two primary families of finite element spaces for meshes of simplices, and the spaces  $\mathcal{Q}_r \mathcal{A}^k$  and  $\mathcal{S}_r \mathcal{A}^k$  on the right side are for meshes of cubes or boxes. Each is defined in any dimension  $n \geq 2$  for each value of the polynomial degree  $r \geq 2$ , and each value of  $d \leq k$ . The parameter  $n$  refers to the operator: the spaces consist of differential  $k$ -forms which belong to the domain of the  $d$ th exterior derivative. Thus for  $n=2$  the spaces describe the Osobolev space<sup>8</sup> of the domain of the curl operator for  $k=1$ , the discrete H(curl) domain of the curl for  $k=2$ , and the discrete H(div) domain of the divergence for  $k=3$ . The spaces  $\mathcal{P}_r \mathcal{A}^k$  and  $\mathcal{P}_r \mathcal{A}^k$  which coincide with the earliest finite elements going back to the case  $n=1$  of linear elements to Courant<sup>9</sup> and collectively referred to as the Lagrange elements. The spaces  $\mathcal{P}_r \mathcal{A}^k$  and  $\mathcal{P}_r \mathcal{A}^k$  which also coincide with the discontinuous Galerkin elements, consisting of piecewise polynomials with no interelement continuity imposed. First introduced by Reed and Milne<sup>10</sup> the space  $\mathcal{P}_r \mathcal{A}^k$  of 2-dim elements was introduced by Gassner and Thurner<sup>11</sup> and generalized to the 3-dimensional spaces  $\mathcal{P}_r \mathcal{A}^k$  and  $\mathcal{P}_r \mathcal{A}^k$  by Arnold, Brezzi, and Fortin<sup>12</sup> due to Brezzi, Douglas and Marini<sup>13</sup> in 2-dimensions. Its generalization to 3-dimensions goes due to Arnold, Brezzi, and Fortin<sup>14</sup>. The undifferentiated and nodal  $\mathcal{P}_r \mathcal{A}^k$  and  $\mathcal{P}_r \mathcal{A}^k$  families are due to Arnold, Falk and Wheeler<sup>15</sup> as part of finite element exterior calculus, extending earlier work of Nipoti<sup>16</sup> for the  $\mathcal{P}_r \mathcal{A}^k$  family.<sup>17</sup> The space  $\mathcal{P}_r \mathcal{A}^k$  is the span of the elementary forms introduced by Whitney.<sup>18</sup>

The family  $\mathcal{Q}_r \mathcal{A}^k$  of cubical elements can be derived from the 1-dim element Lagrange and discontinuous Galerkin elements by a tensor product construction detailed by Arnold, Buffi and Bruneau,<sup>19</sup> but for the most part were presented individually along with the corresponding simplicial elements in the papers mentioned. The second cubical family  $\mathcal{S}_r \mathcal{A}^k$  is due to Arnold and Brezzi.<sup>20</sup>

The finite elements in this table have been implemented as part of the FEM-COSMO project.<sup>21-23</sup> Each may be referenced in the Unified Form Language (UFL) by giving its family, shape, and degree, with the family as observed in the table. For example, the space  $\mathcal{P}_1 \mathcal{A}^1(\Delta_3)$  may be referred to in UFL as: `P1A1T3`.  
 For the  $\mathcal{P}_r \mathcal{A}^k$  family: `P(r)A(k)T(n)`, tetrahedron, 3.  
 For the  $\mathcal{P}_r \mathcal{A}^k$  family: `P(r)A(k)T(n)`, tetrahedron, 3.  
 For the  $\mathcal{Q}_r \mathcal{A}^k$  family: `Q(r)A(k)T(n)`, cube, n.  
 For the  $\mathcal{S}_r \mathcal{A}^k$  family: `S(r)A(k)T(n)`, shape, r, k.  
 For the  $\mathcal{P}_r \mathcal{A}^k$  family: `P(r)A(k)T(n)`, shape, r, k.  
 For the  $\mathcal{P}_r \mathcal{A}^k$  family: `P(r)A(k)T(n)`, shape, r, k.  
 For the  $\mathcal{P}_r \mathcal{A}^k$  family: `P(r)A(k)T(n)`, shape, r, k.

### References

- R. Courant, *Bulletin of the American Mathematical Society* 48, 1943.
- W. H. Reed and T. R. Hill, *Los Alamos Report LA-23-492*, 1973.
- F.A. Buzard and J.M. Thomas, *Acta Numerica* 606, Springer, 1997.
- J.C. Nédélec, *Numerische Mathematik* 35, 1980.
- R. Brezzi, J. Douglas, Jr., and G. Marini, *Numerische Mathematik* 42, 1985.
- D.N. Arnold, R.E. Falk, and R. Winther, *Acta Numerica* 15, 2006.
- H. Harizan, *Mathematics of Computation* 68, 1999.
- H. Winther, *Geometric Integration Theory*, 2007.
- D.N. Arnold, D. Buffi, and G. Sangalli, *Numerische Mathematik*, 2014.
- D.N. Arnold and G. Awanou, *Mathematics of Computation* 2013.
- A. Logg, K.A. Mardal, and G.N. Wells (eds.), *Automated Solution of Differential Equations by the Finite Element Method*, Springer, 2012.
- A. Logg, K.A. Mardal, and G.N. Wells (eds.), *Automated Solution of Differential Equations by the Finite Element Method*, Springer, 2012.
- D.N. Arnold, R.E. Falk, and G.N. Wells (eds.), *Automated Solution of Differential Equations by the Finite Element Method*, Springer, 2012.
- A. Logg and G.N. Wells, *ACM Transactions on Mathematical Software* 27, 2010.
- M.E. Hoot, M.E. Hoot, and G.N. Wells, *ACM Transactions on Mathematical Software* 40, 2014.

